



Effective Oxidation of Alcohols with H_5IO_6 Catalyzed by Nickel(II) Schiff Base Complexes

Dileep Ramakrishna & Badekai Ramachandra Bhat

To cite this article: Dileep Ramakrishna & Badekai Ramachandra Bhat (2010) Effective Oxidation of Alcohols with H_5IO_6 Catalyzed by Nickel(II) Schiff Base Complexes, *Synthesis and Reactivity in Inorganic, Metal-Organic, and Nano-Metal Chemistry*, 40:8, 516-520, DOI: [10.1080/15533174.2010.503180](https://doi.org/10.1080/15533174.2010.503180)

To link to this article: <https://doi.org/10.1080/15533174.2010.503180>



Published online: 31 Aug 2010.



Submit your article to this journal [↗](#)



Article views: 79



View related articles [↗](#)

Effective Oxidation of Alcohols with H_5IO_6 Catalyzed by Nickel(II) Schiff Base Complexes

Dileep Ramakrishna and Badekai Ramachandra Bhat

Catalysis and Materials Chemistry Division, Department of Chemistry, National Institute of Technology Karnataka, Surathkal, Srinivasanagar, India

Nickel(II)-Schiff base-triphenylphosphine complexes catalyze oxidation of alcohols to carbonyls in presence of minimum amount of periodic acid. The catalytic oxidation was developed in mild conditions and showed good yields. The effects of temperature, time, and concentrations of catalyst and co-oxidant were studied. Higher catalytic activity has been observed for NiL1 compared to the other complexes.

Keywords Ni(II) complex, oxidation, periodic acid, Schiff base

INTRODUCTION

The oxidation of alcohols to the corresponding aldehydes and ketones is a key reaction in synthetic organic chemistry, and numerous oxidizing reagents are available to affect this important reaction.^[1–4] Many of these oxidation systems require stoichiometric amounts of metal compounds with concomitant environmental problems. For example, the chromium oxidants are used in vast amounts both in the laboratory and industry.^[5] Recent efforts have focused on systems that employ metal compounds only in catalytic amounts together with a stoichiometric oxidant.^[6–9] Several reaction conditions have been developed for the oxidation of alcohols to carbonyls, such as $RuCl_3/H_5IO_6$,^[10] RuO_4 ,^[11] $RuCl_3/K_2S_2O_8$,^[12] $Ru-Co$ bimetallic catalyzed oxidation,^[13,14] $Cu(MnO_4)_2$ catalyzed oxidation,^[15] TEMPO (2,2,6,6-tetramethylpiperidinyl-1-oxy)-catalyzed oxidation with sodium hypochlorite,^[16–21] Na_2WO_4/H_2O_2 ,^[22] and $Cu(II)$ -salen complex/ H_2O_2 .^[23] The use of transition-metal complexes as catalysts for oxidation reactions has received good attention during the last decades, particularly by the interest in understanding reactions where the metal ion plays a central role in the catalytic reaction. However, despite the extensive studies carried out with cobalt, iron and manganese complexes, only a few investigations have focused on nickel catalysts.^[24–26]

Very recently, we developed an efficient method for the selective oxidation of alcohols to the corresponding aldehydes and ketones using nickel(II) complex containing triphenylphosphine and a Schiff base as a catalyst in ionic liquid media.^[27] As a part of our ongoing research towards nickel catalyzed oxidation chemistry, herein, we report the catalytic oxidation of benzylic and allylic alcohols to the corresponding carbonyl compounds using periodic acid at ambient temperature. This procedure is very simple, mild and clean and works efficiently without any additives.

EXPERIMENTAL

The complexes were synthesized according to a reported literature procedure,^[27] as shown in Scheme 1.

Catalytic Experiment

A typical experimental procedure was as follows: A solution of nickel complex (11 mg, 0.02 mmol) in 20 cm³ CH_3CN was added to the solution of substrate (1 mmol) and H_5IO_6 (230 mg, 1 mmol). The reaction mixture was stirred at 70°C for 30 min and the mixture was extracted with ether. The ether solution was analyzed by GC. Products were isolated by distillation.

Product Analysis

The reaction product analysis was carried out using gas chromatography (GC) (Shimadzu 2014, Japan); the instrument has a 5% diphenyl and 95% dimethyl siloxane Restek capillary column (30 m length and 0.25 mm diameter) and a flame ionization detector (FID). Nitrogen gas was used as the carrier gas. The retention times for different compounds were determined by injecting commercially available compounds under identical gas chromatography conditions. The oxidation products are commercially available, and were identified by GC co-injection with authentic samples.


RESULTS AND DISCUSSION

Catalytic Oxidation

Benzylalcohol was selected as a model substrate for optimization of the process. In order to study the effect of time

Received 20 March 2010; accepted 19 May 2010.

Address correspondence to Badekai Ramachandra Bhat, Catalysis and Materials Chemistry Division, Department of Chemistry, National Institute of Technology Karnataka, Surathkal, Srinivasanagar 575025, India. E-mail: chandpoorna@yahoo.com



NiL1: R=H, NiL2: R= Cl, NiL3: R=Br, NiL4: R=NO₂, NiL5: R= OCH₃.

SCH. 1. Synthesis of nickel (II) complexes.

on the activity, the product analysis was done at regular intervals of time under similar reaction conditions (Figure 1). It was observed that the total reaction time was only 30 minutes. This implies that Ni(II)-complex/CH₃CN-H₅IO₆ catalytic system showed good efficiency. Therefore this catalytic system was studied in detail (Table 2). The oxidation occurred only in poor yield by simply bubbling molecular oxygen through reaction mixture under similar reaction conditions (Table 1, entry 2). To evaluate the catalytic effect of catalyst, the oxidation of benzylalcohol was carried out under similar reaction conditions in the absence of catalyst and no conversion was observed (Table 1, entry 3). Alternatively, the reaction was carried out at room temperature under the same conditions in which the reaction was complete after 210 min (Table 1, entry 4). Further, when the oxidation of benzylalcohol was carried out using less amount of catalyst, the yield was low (Table 1, entry 5). Also the reaction was carried out using low amount of oxidant under same conditions and was observed that the yield was low (Table 1, entry 6).




FIG. 1. Effect of time with % conversion.

The oxidation of other benzylic and allylic alcohols was then examined using the optimized reaction conditions. The results for the oxidation of a variety of alcohols are summarized in Table 2. Benzyl alcohols with functional groups were also oxidized smoothly to the corresponding aldehydes in high yields. As expected, secondary alcohols were oxidized more quickly than primary alcohols. Most oxidations of secondary alcohols were completed within 45 min and gave the corresponding ketones cleanly. Oxidation of allylic alcohols to α , β -unsaturated carbonyl compounds has long been of interest. It was found that cinnamaldehyde was obtained from cinnamyl alcohol in $\sim 90\%$ conversion after 30 min of stirring whereas NaIO₄/2,2,6,6-tetramethylpiperidine-1-oxyl/NaBr system gives only 18% of aldehyde from the same substrate.^[28] Interestingly, catalytic property of nickel(II) complexes is not affected even for chelating substrate (*o*-hydroxy benzylalcohol) as it is in many homogeneous catalytic systems. This present catalytic system requires a very short reaction period of about 30–45 minutes compared to reported catalytic systems.^[29]

TABLE 1
Oxidation of benzylalcohol to benzaldehyde^a

Entry	Reaction time (min)	% Conversion
1	30	92.5
2 ^b	300	<20
3 ^c	300	NR ^d
4 ^e	210	90.6
5 ^f	30	56.9
6 ^g	30	64.9

^aUnless otherwise indicated, all reactions were carried out with 1.08g (1 mmol) of benzylalcohol with 11 mg (0.02 mmol) of Ni(II) catalyst 230 mg (1 mmol) of H₅IO₆, stirring at 70°C.

^bThe reaction was carried out under an atmosphere of O₂ instead of H₅IO₆.

^cThe reaction was carried out without Ni(II) catalyst.



















^dNR = reaction did not occur.

^eThe reaction was carried out at room temperature.

^fThe concentration of Ni(II) catalyst was decreased to 0.01 mmol.











^gThe concentration of the oxidant was decreased to 0.5 mmol.

TABLE 2
Oxidation of alcohols catalyzed by Ni(II) complexes^a in CH₃CN-H₅IO₆ system

Alcohols	Product	Time (min)	% Yield of carbonyl compound ^b				
			NiL1	NiL2	NiL3	NiL4	NiL5
		30	92.6	89.5	84.6	85.4	81.0
		30	91.7	86.2	86.9	85.6	80.1
		30	96.4	95.6	84.6	84.6	80.6
		30	90.1	86.2	82.4	88.6	78.4
		40	91.9	85.4	80.1	80.2	80.0
		40	95.6	87.8	89.6	85.2	82.1
		40	92.1	87.2	90.0	80.1	76.3
		35	85.1	90.0	88.3	88.5	79.8
		30	87.1	89.5	82.6	86.1	81.3

(continued on next page)

TABLE 2
Oxidation of alcohols catalyzed by Ni(II) complexes^a in CH₃CN-H₅IO₆ system (continued)


Alcohols	Product	Time (min)	% Yield of carbonyl compound ^b				
			NiL1	NiL2	NiL3	NiL4	NiL5
		30	95.7	91.3	77.6	71.9	80.1
		30	76.2	75.5	76.0	73.6	71.2
		40	80.2	76.6	74.6	78.1	76.5
		30	75.7	74.9	72.6	73.1	72.5
		30	72.1	72.6	71.2	76.1	70.0

^a 1 mmol alcohol, 1 mmol H₅IO₆, 0.02 mmol Ni (II) complex, 20 mL CH₃CN, stirring at 70°C.

^b GC yield, average of three trials.

Spectral evidence for the complex formation between catalyst and substrate was obtained from UV-Vis spectra of catalyst, benzylalcohol and both in CH₃CN. A bathochromic shift of about 15 nm from 350 nm to 365 nm in the spectra of Ni(II)

was observed and a hyperchromicity was observed at 365 nm (Figure 2). This may be attributed to the formation of a Ni-substrate complex.^[30] On the basis of above discussion, a plausible reaction mechanism is given (Scheme 2).



SCH. 2. Proposed reaction mechanism for the catalytic oxidation of alcohols.




FIG. 2. UV-Vis spectral scans of (a) NiL1 + H₅IO₆ in CH₃CN; (b) NiL1 + H₅IO₆ + Benzylalcohol in CH₃CN.

CONCLUSIONS

In conclusion, the results obtained in this research show that the above square planar nickel (II) complexes are excellent catalysts for the simple, selective and very fast oxidation of alcohols to the corresponding carbonyl compounds with periodic acid in CH₃CN. The system works very efficiently in mild conditions.

REFERENCES

1. Hudlicky, M. *Oxidation in Organic Chemistry*, American Chemical Society: Washington, DC, **1990**, pp. 114.

2. Mirafzal, G. A., and Lozera, M. *Tetrahedron Lett.*, **1998**, 39, 7263.
3. Corey, E. J., and Kim, C. U. *J. Am. Chem. Soc.*, **1972**, 94, 7586.
4. Lou, J. D., and Lou, W. X. *Synth. Commun.* **1997**, 27, 3697.
5. Trost, B. M., and Fleming, I. *Comprehensive Organic Synthesis*, Vol. 7 (Oxidation), Eds., Pergamon Press: UK, **1991**.
6. Zondervan, C., Hage, R., and Feringa, B. L. *Chem. Commun.*, **1997**, 419.
7. Coleman, K. S., Lorber, C. Y., and Osborn, J. A. *Eur. J. Inorg. Chem.*, **1998**, 1673.
8. Moody, C. J., and Palmer, F. N. *Tetrahedron Lett.*, **2002**, 43, 139.
9. Dijkman, A., Marino-Gonzales, A., Payeras, A. M. I., Arends, I. W. C. E., and Sheldon, R. A. *J. Am. Chem. Soc.*, **2001**, 123, 6826.
10. Carlsen, P. H. J., Katsuki, T., Martin, V. S., and Sharpless, K. B. *J. Org. Chem.*, **1981**, 46, 3936.
11. Berkowitz, L. M., and Rylander, P. N. *J. Am. Chem. Soc.*, **1958**, 80, 6682.
12. Varma, R. S., and Hogan, M. E. *Tetrahedron Lett.*, **1992**, 33, 7719.
13. Murahashi, S. I., Naota, T., and Hirai, N. *J. Org. Chem.*, **1993**, 58, 7318.
14. Hongbing, J., Mizugaki, T., Ebitani, K., and Kaneda, K. *Tetrahedron Lett.*, **2002**, 43, 7179.
15. Noureldin, N. A., and Donald, G. L. *J. Org. Chem.*, **1982**, 47, 2790.
16. Nooy, A. E., de J., Besemer, A. C., and Bekkum, H. V. *Synthesis*, **1996**, 1153.
17. Anelli, P. L., Biffi, C., Montanari, F., and Quici, S. *J. Org. Chem.*, **1987**, 52, 2559.
18. Tashino, Y., and Togo, H. *Synlett*, **2004**, 2010.
19. Miyazawa, T., Endo, T., Shiihashi, S., and Okawara, M. *J. Org. Chem.*, **1985**, 50, 1332.
20. Rychnovsky, S. D., and Vaidyanathan, R. *J. Org. Chem.*, **1999**, 64, 310.
21. Zhao, M., Li, J., Song, Z., Desmond, R., Tschaen, D. M., Grabowski, E. J. J., and Reider, P. J. *J. Org. Chem.* **1999**, 64, 2564.
22. Sato, K., Aoki, M., Tagaki, J., and Noyori, R. *J. Am. Chem. Soc.*, **1997**, 119, 12386.
23. Velusamy, S., and Punniyamurthy, T. *Eur. J. Org. Chem.* **2003**, 3913.
24. Masoud, S-N., and Ahmad, A. *Applied Catalysis A*, **2005**, 46, 290.
25. Zhou, W., Hu, B., and Liu, Z. *Applied Catalysis A*, **2009**, 136, 358.
26. Wu, Z.-Y., Xu, D.-J., and Feng, Z.-X. *Polyhedron*, **2001**, 20, 281.
27. Ramakrishna, D., Ramachandra Bhat, B., and Karvembu, R. *Catal. Commun* **2010**, 11, 498.
28. Lei, W., Hu, R.-J., and Wang, Y.-G. *Tetrahedron* **2006**, 62, 8928.
29. Xu, L., and Trudell, M. L. *Tetrahedron Lett.* **2003**, 44, 2553.
30. Jordan, R. B. *Reaction Mechanisms of Inorganic and Organometallic Systems*, 3rd Edition, Oxford University Press: New York, **2007**.